**Directional Infinity on the Riemann Sphere: A Projective-Limit Calculus for Vanishing Complex Exponentials**

Douglas C. Youvan

doug@youvan.com

www.youvan.ai

February 3, 2026

The complex exponential exp(z) never equals zero for any finite complex input z, a fact that is as basic as it is constraining. Yet in analysis and geometry we constantly speak as if functions "vanish at infinity," using boundary language to compress limit behavior into a single symbolic statement. This paper formalizes that compression for the specific case exp(i pi theta n), where the oscillatory nature of imaginary exponents appears to forbid decay. The key move is to treat infinity not as a number but as an ideal boundary point on the Riemann sphere (or equivalently the complex projective line), and to attach to that point an explicit approach ray. For nonzero theta we introduce the ray gamma_theta(t) = -t / (i pi theta), along which the exponent becomes real and tends to minus infinity, forcing exp(i pi theta n) to converge to zero in a clean directional sense. We then develop a lightweight "directional evaluation at infinity" notation f(n_infty; gamma) and show how it unifies asymptotic statements, boundary conditions, and coordinate changes such as w = 1/n that move infinity to a finite puncture. The result is a compact, rigorous framework for discussing vanishing behavior that respects both complex analysis and projective geometry.

Keywords: Riemann sphere, complex projective line, CP1, point at infinity, directional limit, asymptotics, complex exponential, vanishing at infinity, projective compactification, ray to infinity, Möbius transformation, essential singularity, steepest descent, Stokes sectors, boundary functional, limit evaluation.

A collaboration with GPT-5.2-Thinking. CC4.0. 40 pages.

**Outline**

1. Motivation and the Apparent Paradox
1.1 The nonvanishing theorem for the complex exponential
1.2 Why exp(i pi theta n) looks purely oscillatory for real theta and n
1.3 What mathematicians mean by “vanishing at infinity”
1.4 The goal: a precise symbol for decay without contradicting exp(z) != 0

2. Geometry of Infinity
2.1 The Riemann sphere as C union {infty}
2.2 CP1 and projective coordinates as a compactification
2.3 The single point at infinity versus directional approach data
2.4 Why a ray (or path) is part of the definition, not an afterthought

3. Defining n_infty and gamma_theta
3.1 The ideal point n_infty and what it is not
3.2 The ray gamma_theta(t) = -t / (i pi theta) for theta != 0
3.3 Directional evaluation: f(n_infty; gamma) as a boundary functional
3.4 Notational discipline: when exp(i pi theta n_infty) is legitimate shorthand

4. Core Result: Exponential Vanishing as a Directional Limit
4.1 Statement of the lemma: exp(i pi theta gamma_theta(t)) -> 0
4.2 Proof by reduction to exp(-t)
4.3 Dependence on theta and the theta = 0 obstruction
4.4 What changes if theta is real, complex, or purely imaginary

5. The Direction Matters
5.1 Multiple rays to the same n_infty, different limits for the same function
5.2 Decay, growth, and oscillation via Re(i pi theta n) along a direction
5.3 Sector language: decay regions and boundary rays
5.4 When no direction yields decay: the real-theta oscillatory case and its resolution

6. Coordinate Changes: Sending Infinity to a Finite Puncture
6.1 The Möbius map w = 1/n and the image of gamma_theta
6.2 Interpreting n -> n_infty as w -> 0 along a ray
6.3 exp(i pi theta n) as exp(i pi theta / w): a punctured neighborhood viewpoint
6.4 Directional limits near a puncture and why “approach direction” persists

7. A Minimal Calculus of Directional Infinity
7.1 Existence criteria for f(n_infty; gamma) for common function classes
7.2 Rational functions: poles, finite limits, and divergence at n_infty

7.3 Exponentials and exponential polynomials: dominance by real parts
7.4 Warnings: why naive algebra with n_infty produces contradictions

8. Applications as Boundary Conditions and Selection Rules
8.1 Using f(n) -> 0 as n -> n_infty along gamma_theta to pick solutions
8.2 A template for ODE and recurrence problems with “decay at infinity” conditions
8.3 Regularization intuition: damping as a geometric choice of direction
8.4 Relation to steepest descent contours and “best” decay directions

9. Extensions and Variants
9.1 Families of rays gamma(theta, phi) and directional parameterization
9.2 Encoding direction intrinsically: blow-up of infinity and a boundary circle of directions
9.3 Beyond exp(i pi theta n): general exp(a n) and exp(p(n)) cases
9.4 Multivariable generalization: infinity in C^k and approach cones

10. Conclusion
10.1 The resolution: no contradiction with exp(z) != 0
10.2 Infinity as geometry plus approach data, not a magical value
10.3 A compact language for vanishing, growth, and oscillation at the boundary
10.4 What the framework buys: rigor, reuse, and clean asymptotic reasoning

11. References

Art

## 1. Motivation and the Apparent Paradox

The starting point is a tension that shows up whenever we try to speak compactly about limits in complex analysis. On the one hand, the complex exponential is famously well-behaved: it is entire, never hits zero, and carries a rigid multiplicative structure that makes it one of the cleanest functions in mathematics. On the other hand, asymptotic language routinely compresses behavior "far away" into shorthand statements like "this term vanishes at infinity," or "the solution decays at infinity," even when the function under discussion is built from exponentials. The paradox becomes especially sharp for exp(i pi theta n): if theta and n are real, the exponent is purely imaginary, so the value lives on the unit circle. It seems impossible for such a quantity to approach zero. Yet the machinery of complex geometry suggests a way to reconcile the two viewpoints: infinity is not a number, but a boundary point; and the path of approach to that boundary carries the analytic content. The work of this paper is to make that reconciliation explicit and reusable, so that "vanishing" becomes a rigorously defined directional limit at an ideal point, not a contradictory claim about the range of exp.

### 1.1 The nonvanishing theorem for the complex exponential

A foundational fact is that exp(z) is never zero for any complex z. There are several equivalent ways to see this, and each highlights a different aspect of why "forcing exp to be zero" is not just difficult, but logically incompatible with its defining structure.

The cleanest argument comes from the group property. The exponential satisfies exp(z) exp(-z) = exp(0) = 1 for all complex z. This means every value exp(z) has a multiplicative inverse, namely exp(-z). But zero has no multiplicative inverse in any field or integral domain. Therefore exp(z) cannot be zero for any z in the complex numbers.

A second viewpoint uses the exponential's polar form when the input is imaginary. For real x, exp(i x) = cos(x) + i sin(x), whose magnitude is 1. If exp(i x) had magnitude 0 at some x, that would force cos(x) and sin(x) to both be 0, which is impossible. That argument only covers purely imaginary inputs, but it is psychologically useful because it matches the everyday "unit circle" intuition.

A third viewpoint is analytic: exp is an entire function with no zeros, so 1/exp is also entire. This is a strong structural statement: exp is not just nonzero pointwise; it is globally nonvanishing in a way that interacts with complex differentiation and analytic continuation. In all three framings, the conclusion is the same: if we remain in ordinary complex analysis with the usual definition of exp, the equation exp(z) = 0 has no solutions, and any attempt to invent one by algebraic manipulation creates contradictions.

### 1.2 Why exp(i pi theta n) looks purely oscillatory for real theta and n

The expression exp(i pi theta n) appears maximally resistant to decay because the factor i forces the exponent to lie on the imaginary axis whenever theta n is real. In that common situation, exp(i pi theta n) sits on the unit circle for all n, and its magnitude is exactly 1. So as n grows large along the real line, the value does not shrink; it keeps rotating. It may repeat periodically if theta is rational, or wrap densely if theta is irrational, but in either case it does not converge to 0.

This is the intuition that makes the "paradox" feel genuine. If a function's magnitude is identically 1, how could any meaningful notion of "vanishing" apply? The answer is that the apparent impossibility is tied to an implicit restriction: we are assuming n goes to infinity along a particular path, usually the real axis, and we are assuming theta stays real. Both are strong geometric choices. Once we allow n to approach infinity through the complex plane and we allow the approach direction to be part of the statement, the exponent i pi theta n is no longer constrained to remain purely imaginary. Its real part can become negative and unbounded along suitable rays. Along those rays, the exponential does exactly what exponentials do: it decays.

So the "purely oscillatory" appearance is not wrong; it is simply incomplete. It describes the behavior on one slice of the domain (real n with real theta). The broader complex-analytic picture asks a different question: among all possible ways to go to infinity in the complex plane, which ones make the real part of i pi theta n go to minus infinity? Those are the directions where decay becomes available. This shift of viewpoint is the doorway to a rigorous resolution.

### 1.3 What mathematicians mean by "vanishing at infinity"

In practice, "vanishing at infinity" is a compact phrase that stands in for a limit statement. For functions on the real line, it typically means lim(x -> +infty) f(x) = 0, or sometimes lim(|x| -> infty) f(x) = 0. But in complex analysis the phrase is more delicate because the complex plane has many directions to infinity, and different directions can produce different asymptotic behavior.

There are several standard ways to formalize the idea, and the differences matter.

One approach uses the one-point compactification of the complex plane, the Riemann sphere. Under this compactification, infinity is a single added point, and a function "vanishes at infinity" if it extends continuously to that point with value 0. In plain language: f vanishes at infinity if f(n) -> 0 whenever n -> infinity, no matter how you go to infinity. This is a strong notion; many functions do not satisfy it because different directions produce different limits or no limit at all.

A second approach is directional: you specify a path or ray gamma(t) that escapes to infinity, and you call f "vanishing at infinity along gamma" if lim(t -> +infty) f(gamma(t)) = 0. This is a weaker notion, but it is the one that actually matches most asymptotic reasoning in applied

complex analysis, steepest descent, and boundary-value problems. The direction is not an incidental detail; it is the entire content.

A third approach uses neighborhoods of infinity via a change of variable w = 1/n. Then “n near infinity” corresponds to “w near 0.” Under this dictionary, “vanishing at infinity” becomes “vanishing near w = 0” along whatever approach in w corresponds to the chosen approach in n. This method is especially useful because it translates a boundary-at-infinity problem into a local analysis near a point, while still preserving directional information.

For this paper’s purposes, the directional notion is the most honest: it captures exactly what we mean when we say exp(i pi theta n) can “vanish at infinity” even though it never equals 0 for any finite n. It vanishes in the sense of a limit along a specific escape path.

**1.4 The goal: a precise symbol for decay without contradicting exp(z) != 0**

The goal is not to manufacture a forbidden solution to exp(z) = 0, and not to redefine the exponential. The goal is to name, precisely and compactly, a boundary phenomenon that people already use informally: an expression can tend to 0 as its input tends to infinity along a chosen direction.

To do that, we introduce a disciplined notation that separates three ideas that are often conflated:

1. The ideal point n_infty, which is the point at infinity on the Riemann sphere.
2. The approach path gamma_theta, which records how n goes to infinity.
3. The directional limit functional f(n_infty; gamma_theta) defined as lim(t -> +infty) f(gamma_theta(t)) when it exists.

With these in place, the statement “exp(i pi theta n) vanishes at infinity” becomes a precise sentence: along the ray gamma_theta(t) = -t / (i pi theta), the real part of the exponent equals -t, so exp(i pi theta gamma_theta(t)) = exp(-t) and therefore the limit is 0. The exponential never equals 0 at any finite input, and the nonvanishing theorem remains untouched. What changes is the grammar: instead of asking for a value of n that makes the exponential equal to 0, we define a boundary evaluation at n_infty along a prescribed ray, and we show that this boundary evaluation equals 0.

In that sense, n_infty is not a “solution,” and gamma_theta is not a trick. Together they provide a mathematically standard way to say something that is intuitively clear: exponentials decay when the real part of the exponent goes to minus infinity, and in the complex plane you can choose a direction to make that happen.

## 2. Geometry of Infinity

To speak cleanly about “n goes to infinity” in complex analysis, we need a geometry in which infinity is not a vague phrase but an actual location. The point is not to pretend infinity is a number. The point is to treat infinity as a boundary point of a compact space that contains the complex plane as an open dense subset. Once we do that, statements about limits “at infinity” become ordinary statements about limits at a point. This is the structural move that makes asymptotic shorthand mathematically legitimate. However, complex analysis immediately adds a second lesson: even though infinity is a single point in the compactification, the analytic behavior near that point typically depends on how we approach it. The geometry supplies the stage; the path supplies the meaning. This section sets up the minimal geometric language needed to make n_infty and gamma_theta feel natural rather than ad hoc.

### 2.1 The Riemann sphere as C union {infty}

The Riemann sphere is the most familiar compactification of the complex plane. It is the set C union {infty} equipped with a topology that makes it homeomorphic to the 2-sphere. Geometrically, one imagines the complex plane sitting tangent to a sphere at the south pole, and then uses stereographic projection to identify each complex number with a point on the sphere. The north pole corresponds to the added point infinity. In this model, “n goes to infinity” simply means “the corresponding point on the sphere approaches the north pole.”

This is already a conceptual upgrade: instead of a statement about escaping all bounded regions in C, we can speak about convergence to a point in a compact space. The topology is chosen precisely so that sequences with |n_k| -> infinity converge to the point infinity on the sphere. In this setting, a function f on C “has a limit at infinity” if it extends continuously to the north pole. And “f vanishes at infinity” means that extension takes the value 0 at the north pole.

The strength of the Riemann sphere model is that it is canonical and compatible with the analytic structure of complex functions. Many standard operations in complex analysis become cleaner when one allows infinity as a point. For instance, rational functions can be understood globally as meromorphic maps on the sphere, with their poles and zeros treated symmetrically. Infinity itself behaves like a pole or zero depending on the degree balance in the rational function. This symmetry is one reason infinity feels so natural in complex analysis: it is not an anomaly; it is a point with local analytic meaning.

However, the same compactness that makes the Riemann sphere elegant also forces a caution: there is only one added point. Different directions in the plane all converge to that same point. If you want to preserve directional information, you must encode it separately, because the topology alone does not distinguish approach angles.

**2.2 CP1 and projective coordinates as a compactification**

The same object can be described algebraically as the complex projective line, CP1. This viewpoint is often the cleanest for writing: CP1 is the set of equivalence classes [z:w] of nonzero pairs (z,w) in C^2 under the relation (z,w) ~ (lambda z, lambda w) for any nonzero complex lambda. Think of [z:w] as homogeneous coordinates. The complex plane embeds into CP1 by sending a complex number n to [n:1]. The missing point corresponds to [1:0], which is the projective way to write infinity.

This is not just a change of costume. Projective coordinates explain why adding one point is enough to compactify the plane: CP1 is already compact as a complex manifold. The map n -> [n:1] identifies C with CP1 minus the single point [1:0]. So the “compactification” is literally the act of adding back that one missing point.

The projective viewpoint also prepares the ground for transformations that move infinity around. Möbius transformations are precisely the automorphisms of CP1. They act transitively on the sphere, meaning any point can be sent to any other point by an appropriate Möbius map. In affine coordinates, a typical Möbius map looks like (a n + b)/(c n + d), but in projective coordinates it becomes a linear action on [z:w] via a 2x2 matrix. This makes infinity a fully legitimate point in the same space as finite numbers: it transforms just as naturally as any other point.

For our purposes, CP1 provides a neat language for the change of variable w = 1/n. In projective terms, w = 1/n is simply the coordinate change between the chart w = z/w and the chart n = z/w, each chart valid away from one point. Infinity is not a special metaphysical place; it is the point where one chart fails and the other becomes appropriate.

**2.3 The single point at infinity versus directional approach data**

Here is the subtlety that matters for exp(i pi theta n). On CP1 there is exactly one point at infinity. If you take any ray n(t) = t e^(i phi) with t -> +infty, it converges to that same point [1:0]. In the topology of the Riemann sphere, all ways of escaping to infinity look the same at the level of the limit point. This is wonderful for compactness and global function theory, but it also means the sphere does not remember how you got there.

Analytically, that missing information can be crucial. Many functions do not have a single well-defined limit at infinity that is independent of direction. Even when a function is bounded along one ray, it may blow up along another. Even when it tends to a limit along one ray, it may oscillate or fail to converge along another. So the single point infinity on the sphere is sometimes too coarse as a descriptor of asymptotic behavior.

This is why our notation separates n_infty from gamma. The symbol n_infty names the boundary point in CP1. The symbol gamma_theta names the approach path. The limit

functional f(n_infty; gamma_theta) is not the same as the unconditional limit at infinity. It is a directional limit. The approach path carries the asymptotic content.

In fact, this resolves the earlier tension in a clean way. For real theta and real n along the real axis, exp(i pi theta n) has magnitude 1 and does not decay. But if we allow n to approach infinity along a complex ray chosen to force the real part of i pi theta n to go to minus infinity, then the exponential decays. Both statements are simultaneously true because they refer to different approach paths to the same ideal point.

So infinity is a single point, but asymptotic analysis is often a story about how functions behave as we approach that point from different directions. The point gives a global address; the direction gives the local meaning.

**2.4 Why a ray (or path) is part of the definition, not an afterthought**

It is tempting to treat the path as an informal add-on, something you mention only when needed. But in complex analysis, if you omit the path you frequently change the statement from true to false.

Consider the phrase “f tends to 0 at infinity.” If interpreted as an unconditional limit on the Riemann sphere, it means f extends continuously to n_infty with value 0. For exp(i pi theta n), that unconditional statement is false in general, because along many paths to infinity the function does not approach 0 at all. Along the real axis with real theta, it stays on the unit circle. Along other directions it may blow up or decay depending on the sign of the real part of the exponent.

But if we interpret “tends to 0 at infinity” directionally, then it becomes a statement about a chosen path. That statement can be true, and in fact can be engineered by choosing the path so that the exponent’s real part becomes large and negative. This is not a trick; it is exactly what steepest descent and contour deformation are built to do. One selects curves along which an exponential term decays fastest, making integrals converge and asymptotic expansions legitimate.

So the path is not an aesthetic choice. It is the difference between an unconditional boundary value and a directional boundary value. In our framework, gamma_theta is part of the data of the claim. Writing exp(i pi theta n_infty; gamma_theta) = 0 is not merely a more precise version of exp(i pi theta n_infty) = 0; it is a different kind of statement, one that preserves the nonvanishing theorem while still capturing the decay phenomenon that motivated the discussion.

This section’s geometric conclusion is simple: CP1 gives you a place called infinity, but complex analysis requires that you also specify how you approach that place. Infinity provides the boundary point; the ray provides the asymptotic meaning.

**3. Defining n_infty and gamma_theta**

This section introduces the two pieces of data that replace the impossible idea of “solve exp(i pi theta n) = 0 for n.” The first piece is geometric: n_infty names the ideal point at infinity in the compactified domain. The second piece is analytic: gamma_theta is an explicit ray along which the exponent becomes large and negative in its real part, so the exponential decays. Once those two are in place, we define a boundary functional that evaluates limits at infinity along a specified path. The end result is a small, disciplined calculus for statements that look like “a function vanishes at infinity,” while keeping the nonvanishing theorem for the exponential intact.

**3.1 The ideal point n_infty and what it is not**

We work on the extended complex plane C union {infty}, also known as the Riemann sphere, or equivalently on CP1. The symbol n_infty denotes the point at infinity in this compactification. Its role is purely geometric: it provides a name for the boundary point approached by sequences (or curves) whose modulus tends to infinity.

What n_infty is:

- A point in the compactified space that is the limit of any sequence n_k in C with |n_k| -> infinity.
- A boundary marker that allows us to speak of convergence “at infinity” the same way we speak of convergence at any finite point.

What n_infty is not:

- It is not a complex number.
- It is not an element of the field C in the algebraic sense.
- It is not something you can insert into formulas and then manipulate with ordinary arithmetic without care.

The last point matters. Many familiar expressions become meaningless or ambiguous if we treat infinity as a number. For example, n_infty - n_infty, 1/n_infty, or n_infty * 0 do not have consistent field-like meanings. In the compactification, 1/n is a legitimate function on C minus {0}, but at infinity it is interpreted by taking limits: 1/n tends to 0 as n -> n_infty. That is a limit statement, not an algebraic identity.

So n_infty should be treated as an address on the boundary of the domain, not as a new value in the number system. Whenever we write something involving n_infty, it must ultimately mean a statement about limits in C as n escapes to infinity.

**3.2 The ray gamma_theta(t) = -t / (i pi theta) for theta != 0**

Fix theta with theta != 0. Define a ray gamma_theta : (0, infty) -> C by

gamma_theta(t) = -t / (i pi theta), for t > 0.

This ray is chosen for a single reason: it converts the complex exponent into a purely real, strictly decreasing quantity. Substituting gamma_theta into the exponent gives

i pi theta gamma_theta(t)
= i pi theta ( -t / (i pi theta) )
= -t.

So along this ray, the exponential becomes

exp(i pi theta gamma_theta(t)) = exp(-t),

which tends to 0 as t -> +infty.

Several remarks clarify what this means.

First, gamma_theta is not arbitrary. It is the unique straight ray through the origin that solves the equation

i pi theta n = -t

with t real and positive. In other words, gamma_theta is the steepest decay direction for the exponential exp(i pi theta n) among straight rays parameterized by a positive real t.

Second, gamma_theta depends on theta. If theta changes, the direction in the complex plane along which the exponential decays changes. This dependence is not a nuisance; it is part of the structure. It is exactly the statement that decay is controlled by the sign and direction of the real part of the exponent.

Third, gamma_theta is a path to infinity: |gamma_theta(t)| grows linearly with t. Therefore gamma_theta(t) approaches n_infty in the topology of the Riemann sphere. This is why gamma_theta is the bridge between the geometry of infinity and the analysis of decay.

**3.3 Directional evaluation: f(n_infty; gamma) as a boundary functional**

To make these ideas reusable, we define a simple boundary functional that turns a directional limit into a piece of notation.

Let gamma : (0, infty) -> C be any curve such that |gamma(t)| -> infinity as t -> +infty. Then gamma approaches n_infty. For any function f defined on a set containing the image of gamma for large t, define the directional evaluation of f at n_infty along gamma by

f(n_infty; gamma) = lim(t -> +infty) f(gamma(t)),

whenever this limit exists.

This definition does several things at once:

- It makes “evaluation at infinity” a standard limit.
- It forces us to record the approach path, which is the analytic content.
- It allows us to distinguish between unconditional behavior at infinity and directional behavior along a chosen path.

For example, for f(n) = exp(i pi theta n) and gamma = gamma_theta, we have

f(n_infty; gamma_theta) = lim(t -> +infty) exp(i pi theta gamma_theta(t))
= lim(t -> +infty) exp(-t)
= 0.

So the statement “exp(i pi theta n) vanishes at infinity” becomes a precise claim:

exp(i pi theta n)(n_infty; gamma_theta) = 0.

This avoids any suggestion that exp hits 0 at a finite argument. It is purely a boundary limit statement.

This functional viewpoint also makes clear what can go wrong. If you choose a different gamma, the limit may differ or may not exist. In particular, along the real axis with real theta, the limit does not approach 0. The notation forces you to say which gamma you mean.

**3.4 Notational discipline: when exp(i pi theta n_infty) is legitimate shorthand**

The temptation is to write exp(i pi theta n_infty) = 0. Taken literally, this is not well-formed, because n_infty is not a complex number and exp is initially defined on complex numbers. However, mathematicians often use boundary shorthand when the intended meaning is unambiguous within a defined convention.

In this paper, the shorthand is legitimate only under an explicit rule:

Rule: Any expression of the form f(n_infty) is shorthand for f(n_infty; gamma) with a gamma that has been specified in the current context, or that is fixed by convention for the discussion.

In our setting, if gamma_theta has been declared as the approach path, then writing

exp(i pi theta n_infty) = 0

is acceptable shorthand for

exp(i pi theta n_infty; gamma_theta) = 0,

meaning

lim(t -> +infty) exp(i pi theta gamma_theta(t)) = 0.

But without gamma_theta (or another chosen path), the statement is misleading, because it suggests an unconditional limit at infinity that is generally false.

So the discipline is simple:

- Use exp(i pi theta n_infty; gamma_theta) when you want precision.
- Allow exp(i pi theta n_infty) only after gamma_theta is fixed and the audience is reminded that infinity is being evaluated directionally.

This discipline preserves the nonvanishing theorem: exp(z) never equals 0 for any finite z. What we are doing is not solving exp(z) = 0. We are defining a boundary evaluation at an ideal point and choosing an approach ray that turns oscillation into decay.

**4. Core Result: Exponential Vanishing as a Directional Limit**

This section states and proves the central mathematical fact around which the rest of the framework is organized: although exp(z) never equals zero for any finite z, the composite exp(i pi theta n) can be made to converge to zero as n approaches n_infty along a specifically chosen ray. The proof is deliberately elementary. It is not an analytic continuation trick or a redefinition of exp. It is simply the observation that "imaginary-looking" exponents can be made real and negative by choosing a complex direction for n. Once that is done, the exponential behaves exactly as expected: it decays to 0. We also make explicit what is and is not possible depending on theta, because theta controls the geometry of the decay direction.

**4.1 Statement of the lemma: exp(i pi theta gamma_theta(t)) -> 0**

Fix theta with theta != 0. Define the ray

gamma_theta(t) = -t / (i pi theta), for t > 0.

Then the lemma is:

Lemma. As t -> +infty, exp(i pi theta gamma_theta(t)) -> 0.

Equivalently, in the boundary functional notation introduced earlier, if f(n) = exp(i pi theta n), then

f(n_infty; gamma_theta) = 0.

This is the entire content of the phrase “the exponential vanishes at infinity” in this context. It is not a claim about the value of the exponential at a finite point. It is a claim about a limit taken along a prescribed path that approaches the ideal point n_infty.

**4.2 Proof by reduction to exp(-t)**

The proof is a one-line reduction enabled by the way gamma_theta was chosen.

Starting from the exponent, substitute gamma_theta(t):

i pi theta gamma_theta(t)
= i pi theta ( -t / (i pi theta) )
= -t.

Therefore the composite exponential simplifies exactly to a real exponential:

exp(i pi theta gamma_theta(t)) = exp(-t).

Since exp(-t) -> 0 as t -> +infty, we conclude

exp(i pi theta gamma_theta(t)) -> 0 as t -> +infty.

This completes the proof.

Two points are worth emphasizing.

First, nothing is hidden. The vanishing is not subtle. It is the same vanishing one sees in any real exponential with negative exponent growing in magnitude.

Second, the proof shows why the ray is “the right one.” We did not discover the decay after choosing gamma_theta; we defined gamma_theta precisely so the exponent becomes -t. The ray is effectively an explicit solution to the design requirement “make the exponent real and large negative.”

**4.3 Dependence on theta and the theta = 0 obstruction**

The lemma depends crucially on theta being nonzero. If theta = 0, then the entire expression collapses:

exp(i pi theta n) = exp(0) = 1

for every n. In that case there is no decay direction, no ray gamma_theta, and no meaningful sense in which the function can vanish at n_infty. The obstruction is absolute: it is not a failure of technique; it is a change in the function itself.

When theta != 0, the dependence on theta appears in the direction of gamma_theta. The ray is proportional to 1/theta. If theta is scaled by a nonzero factor, the ray rotates and rescales accordingly. Conceptually, theta determines which directions in the n-plane produce negative

real part in the exponent i pi theta n. The ray gamma_theta is the simplest possible choice that makes the exponent exactly -t, not merely negative.

This also explains an important qualitative point: there is not a single, universal “decay at infinity” for exp(i pi theta n). The decay is direction-dependent and theta-dependent. The only universal statement is that if theta != 0, then there exists at least one ray to infinity along which the exponential decays to 0.

**4.4 What changes if theta is real, complex, or purely imaginary**

The lemma itself does not require theta to be real. It only requires theta != 0. But the interpretation of the result, and the relationship to the “purely oscillatory” intuition, depends on what sort of number theta is.

Theta real.

If theta is real and nonzero, then exp(i pi theta n) is purely oscillatory along the real axis because the exponent is purely imaginary when n is real. That is the origin of the intuition that decay is impossible. The lemma shows that decay becomes possible the moment we allow n to move into the complex plane: the ray gamma_theta points in an imaginary direction (up to sign) and forces the exponent to become real negative. So for real theta, the contrast is sharp: real-axis behavior is oscillation, complex-direction behavior can be decay.

Theta purely imaginary.

Write theta = i b with b real and nonzero. Then the exponent becomes

i pi theta n = i pi (i b) n = -pi b n.

Along the real axis, this is already real. If b > 0, then choosing n -> +infty along the positive real axis makes the exponent tend to -infty and the exponential decays to 0 without any complex detour. If b < 0, the decay happens as n -> -infty along the real axis. So for purely imaginary theta, the situation is almost the opposite of the real-theta case: decay is available on the real axis, and oscillation is not the default.

Theta complex with both real and imaginary parts.

Let theta = a + i b with a and b real. Then

i pi theta n = i pi (a + i b) n = i pi a n - pi b n.

Now the real part of the exponent is controlled by -pi b n when n is real. If b > 0, then along n -> +infty on the real axis you already get decay like exp(-pi b n), modulated by an oscillatory factor from i pi a n. If b = 0, you recover the purely oscillatory real-theta case along the real axis. If b < 0, the real axis produces growth rather than decay, and you must choose a different direction to achieve vanishing.

In the fully complex case, gamma_theta still works exactly as defined. It forces the exponent to equal -t, independent of the internal decomposition of theta. That is its elegance: it is a universal decay ray for any nonzero theta, because it is constructed as the direct inverse of the coefficient i pi theta in the exponent.

Summary of the regimes.

- Real theta: real-axis behavior is oscillatory; decay requires moving n into a complex direction.
- Purely imaginary theta: decay can occur already along the real axis in one direction.
- General complex theta: decay may already happen along the real axis if Im(theta) has the appropriate sign, but gamma_theta guarantees decay for any theta != 0.

This regime analysis reinforces the main conceptual point of the paper: the phrase “vanishing at infinity” is not a statement about an input value. It is a statement about a boundary limit, and in the complex plane boundary limits are inseparable from the direction of approach.

**5. The Direction Matters**

The compactification C union {infty} gives only one point called n_infty, but complex analysis refuses to let that be the end of the story. Infinity is a single location in the Riemann sphere, yet it is approached from infinitely many directions in the plane. For many functions, especially exponentials and exponential polynomials, the limiting behavior as n approaches n_infty is not a property of the point alone but of the path. This is not a pedantic detail. It is the structural reason the phrase “vanishes at infinity” must be handled with care: the same function can decay along one ray, blow up along another, and oscillate along a third, all while heading toward the same ideal point. In the specific problem of exp(i pi theta n), direction is the entire mechanism by which oscillation is turned into decay. This section makes that dependence explicit in a way that can be reused beyond this single example.

**5.1 Multiple rays to the same n_infty, different limits for the same function**

On the Riemann sphere, any curve gamma(t) in C with |gamma(t)| -> infinity converges to the same point n_infty. In that topological sense, all rays are equal: they all land at infinity. But for a function f(n), the limit

lim(t -> +infty) f(gamma(t))

may depend dramatically on gamma. This is the core phenomenon that forces us to treat “evaluation at infinity” as directional unless a stronger, direction-independent limit can be proven.

It is easy to see this with simple examples:

- Take f(n) = n. Along any ray to infinity, f diverges to infinity in magnitude. There is no finite limit at n_infty in any direction, so there is no meaningful “value” at infinity.
- Take f(n) = 1/n. Along any ray to infinity, f tends to 0. Here the limit is direction-independent, and it makes sense to say f vanishes at infinity without specifying gamma.
- Take f(n) = exp(n). Along the positive real axis, exp(n) -> infinity. Along the negative real axis, exp(n) -> 0. Along the imaginary axis, exp(i t) oscillates with magnitude 1 and has no limit. One function, one point n_infty, but three incompatible asymptotic behaviors depending on the ray.

This illustrates why our boundary functional must record the path. Direction-independence is a special property, not the default. The only safe universal statement is: if the limit at infinity exists in a direction-independent sense, then it is the same along every path. But most interesting functions fail that criterion, and exp(i pi theta n) is one of them in the real-theta regime.

**5.2 Decay, growth, and oscillation via Re(i pi theta n) along a direction**

For exponentials, the classification of behavior reduces to a single scalar: the real part of the exponent. Write

exp(i pi theta n) = exp( Re(i pi theta n) ) times a unit-magnitude factor exp( i Im(i pi theta n) ).

The magnitude is

|exp(i pi theta n)| = exp( Re(i pi theta n) ).

So everything comes down to the sign and asymptotics of Re(i pi theta n) along the chosen path.

Let us describe a ray by n = r u, where r > 0 and u is a unit complex number (a direction). As r -> +infty along that ray, we have

Re(i pi theta n) = Re(i pi theta r u) = r Re(i pi theta u).

If Re(i pi theta u) < 0, then Re(i pi theta n) -> -infty and the exponential decays to 0 along that ray.
If Re(i pi theta u) > 0, then Re(i pi theta n) -> +infty and the exponential blows up in magnitude along that ray.
If Re(i pi theta u) = 0, then the magnitude stays constant at 1 and the behavior is purely oscillatory (or at least non-decaying) along that ray.

This is the clean geometric classification:

- Decay directions are those u for which Re(i pi theta u) is negative.
- Growth directions are those u for which Re(i pi theta u) is positive.
- Boundary directions are those u for which Re(i pi theta u) is zero.

The ray gamma_theta introduced earlier is the maximally strict decay ray in the sense that it forces the exponent itself to be exactly -t, not merely negative in real part. It is the direction for which Re(i pi theta u) is as negative as possible among rays normalized by r, because it is chosen so that i pi theta u is a negative real number.

So the “direction matters” statement can be turned into a computable criterion: choose a direction u, evaluate Re(i pi theta u), and you immediately know whether exp(i pi theta n) decays, grows, or oscillates along the ray n = r u.

**5.3 Sector language: decay regions and boundary rays**

In asymptotic analysis, one often describes these direction classes as sectors. A sector is a wedge-shaped region in the complex plane defined by an angular interval. For linear exponentials exp(a n), with a fixed complex constant a, the plane divides into two half-planes of directions: those where Re(a n) is positive and those where it is negative, separated by boundary rays where Re(a n) = 0.

For our case, the constant is a = i pi theta. The set of n for which Re(a n) < 0 is an open half-plane bounded by a line through the origin. Intersecting that half-plane with large-radius circles produces an arc of directions along which decay occurs. The boundary consists of two opposite rays through the origin, defined by the condition Re(a u) = 0, which is equivalent to saying a u is purely imaginary.

This language is useful for two reasons.

First, it replaces an abstract “depends on direction” statement with a geometric picture: there is an entire region of directions that produce decay, not just a single ray. The ray gamma_theta is a convenient representative, but it sits inside a full decay half-plane.

Second, the boundary rays are precisely where qualitative behavior changes. Crossing a boundary ray flips decay into growth or vice versa. In more complicated exponential expressions, such boundary rays generalize into Stokes lines, where dominance relations between competing exponentials switch. Even in this simple linear case, the sector decomposition is the simplest incarnation of that general phenomenon.

So one can summarize the geometry as follows: for theta != 0, the complex plane around the origin has a decay half-plane for exp(i pi theta n), a growth half-plane, and boundary rays separating them. Any path to infinity that eventually stays within the decay half-plane will force

the exponential toward 0. Any path to infinity that eventually stays within the growth half-plane will cause blow-up. Paths that cling to the boundary rays are oscillatory in magnitude.

**5.4 When no direction yields decay: the real-theta oscillatory case and its resolution**
There is a common confusion that is worth addressing directly: if theta is real, exp(i pi theta n) looks like a pure phase factor, so one might conclude there is no direction along which it decays. This confusion is understandable because it usually arises from silently restricting n to be real as well.

If theta is real and n is forced to remain real, then indeed the magnitude is identically 1 and no decay is possible. Along the real axis, Re(i pi theta n) = 0 for all n, so the function is trapped on the unit circle. In that restricted setting, the decay sector collapses to nothing because the only allowed direction is the real direction, which is a boundary direction.

But once n is allowed to be complex, the decay half-plane immediately reappears. For real theta, the constant a = i pi theta is purely imaginary, and the condition Re(a n) < 0 becomes a condition on the imaginary part of n. In fact, writing n = x + i y with x and y real, we have

a n = i pi theta (x + i y) = i pi theta x - pi theta y.

The real part is -pi theta y. If theta > 0, then choosing y > 0 forces the real part negative, producing decay as y grows. If theta < 0, choosing y < 0 produces decay. So for real theta, decay is not prohibited; it is simply located off the real axis, in the upper or lower half-plane depending on the sign of theta.

This is the resolution: “no direction yields decay” is only true if you impose the extra constraint that n must remain real. In the full complex domain, theta != 0 always admits decay directions. The ray gamma_theta is an explicit, canonical choice among them, engineered to make the exponent exactly -t and thus make the decay statement trivial and robust.

So the correct statement is:

- For real theta, exp(i pi theta n) is purely oscillatory along the real axis, but it decays along suitable complex rays.
- The apparent impossibility is an artifact of a restricted domain, not a property of the exponential itself.

This completes the conceptual task of the section: infinity is a single point, but asymptotic behavior is directional; decay, growth, and oscillation are governed by the sign of Re(i pi theta n) along the chosen direction; and the “real-theta oscillation” issue disappears the moment we allow complex approach paths, which is exactly what the geometry of n_infty is designed to sanction._

## 6. Coordinate Changes: Sending Infinity to a Finite Puncture

One of the most powerful habits in complex analysis is to replace “behavior at infinity” with “behavior near a finite point” by a change of coordinates. On the Riemann sphere this is not a trick; it is the natural transition between charts. The map w = 1/n is the simplest Möbius transformation that exchanges 0 and infinity, turning the ideal point n_infty into the ordinary point w = 0. This translation is valuable for two reasons. First, it turns global asymptotic statements into local statements near a point, which can then be analyzed with the familiar machinery of local complex analysis. Second, it makes visible a key fact that can otherwise be missed: direction does not disappear when you move infinity to a finite point. It reappears as the direction of approach to a puncture. In other words, the need to track approach paths is not a quirk of infinity; it is a general feature of local complex behavior near singular or boundary points.

### 6.1 The Möbius map w = 1/n and the image of gamma_theta

The map w = 1/n is a Möbius transformation on the Riemann sphere. In the language of CP1, it swaps the affine charts: n is the coordinate on the chart where [z:w] has w != 0, while w is the coordinate on the chart where [z:w] has z != 0. Under this transformation:

- n = 0 maps to w = infinity.
- n = infinity (that is, n_infty) maps to w = 0.

So w = 1/n is the canonical way to bring infinity into the local neighborhood of a finite point.

Now apply this to our ray gamma_theta(t) = -t / (i pi theta). Its image in the w-plane is

w_theta(t) = 1 / gamma_theta(t)
= 1 / ( -t / (i pi theta) )
= -(i pi theta) / t.

This expression is already illuminating. As t -> +infty, w_theta(t) -> 0. So the path that runs off to infinity in the n-plane becomes a path that runs into 0 in the w-plane. Moreover, it runs into 0 along a specific ray determined by the complex constant -(i pi theta). The magnitude shrinks like 1/t, and the argument is fixed by theta. So gamma_theta has become a straight approach line to the puncture w = 0.

This is a small but important payoff: gamma_theta is no longer something you must accept as a special ray “to infinity.” It is simply the straight-line approach to w = 0 in the reciprocal coordinate. That is, gamma_theta is natural in whichever chart you choose.

**6.2 Interpreting n -> n_infty as w -> 0 along a ray**

The slogan is: n -> n_infty corresponds to w -> 0. More precisely, a sequence n_k in C escapes to infinity if and only if w_k = 1/n_k tends to 0. And a curve gamma(t) with |gamma(t)| -> infinity corresponds to a curve eta(t) = 1/gamma(t) that tends to 0.

However, the directional aspect persists. The n-plane has many ways to approach infinity; the w-plane has many ways to approach 0. The approach angle in the n-plane becomes an approach angle in the w-plane. For our specific ray, w_theta(t) approaches 0 along the direction of -(i pi theta). If theta is real and positive, then -(i pi theta) points straight down the imaginary axis, so w approaches 0 from below along the imaginary direction. If theta has other phases, the approach direction rotates accordingly.

So the statement “n approaches n_infty along gamma_theta” becomes the statement “w approaches 0 along the ray w_theta(t) = -(i pi theta)/t.” This is often the best way to think about the construction: we are not doing something exotic at infinity; we are selecting a ray of approach to a puncture.

This re-framing also helps with intuition. Infinity can feel metaphysical. A punctured neighborhood feels concrete: it is just a small disk around 0 with the center removed, and we are taking a limit as we move along a specified path toward the missing point.

**6.3 exp(i pi theta n) as exp(i pi theta / w): a punctured neighborhood viewpoint**

Under the substitution n = 1/w, our exponential becomes

exp(i pi theta n) = exp(i pi theta / w).

Now the question “what happens as n -> n_infty?” becomes “what happens as w -> 0?” But notice what this does to the function: exp(i pi theta / w) has extremely wild behavior near w = 0. The reason is that 1/w has a pole at 0, and composing exp with a pole produces an essential-type explosion of oscillation and growth/decay depending on direction. Even without naming any classification, the qualitative point is clear: as w gets small, the quantity (i pi theta)/w becomes large, and exponentials of large complex numbers are sensitive to the sign of the real part.

This punctured-neighborhood viewpoint clarifies why an unconditional limit at infinity often fails. If you ask whether exp(i pi theta / w) has a single limit as w -> 0 independent of direction, the answer is generally no. Different approach rays to 0 produce different behaviors because the real part of (i pi theta)/w changes sign across rays.

But along our chosen approach w_theta(t) = -(i pi theta)/t, the expression becomes simple:

(i pi theta) / w_theta(t)
= (i pi theta) / ( -(i pi theta)/t )
= -t.

Therefore

exp(i pi theta / w_theta(t)) = exp(-t) -> 0.

So the core lemma is now a local statement near w = 0: along a particular ray into the puncture, the function decays to 0. This is exactly the same content as before, just expressed in a coordinate system where infinity has been moved into a local neighborhood.

This is also why gamma_theta was "the right ray." In w-coordinates, we literally choose the ray so that (i pi theta)/w is a large negative real number. That is the canonical "steepest decay" choice.

**6.4 Directional limits near a puncture and why "approach direction" persists**

The biggest conceptual lesson of the coordinate change is that direction is not a special complication attached to infinity. It is a universal feature of limits near points where the function has strong directional sensitivity. Infinity is one such point; a puncture like w = 0 for exp(i pi theta / w) is another.

Even in the w-plane, if you approach 0 along different rays, you can force (i pi theta)/w to have real part tending to +infty (blow-up), -infty (decay), or to remain purely imaginary in leading order (unit-magnitude oscillation). This directional dependence is the local version of the sector picture discussed earlier. The boundary between decay and growth is given by rays where Re((i pi theta)/w) = 0 in the limit, which are exactly the rays along which the exponent is asymptotically imaginary.

So the approach direction persists because the function itself couples magnitude to the real part of its exponent, and that real part depends on the angle of approach whenever a pole or infinity is involved. Changing coordinates does not remove the dependence; it simply relocates it.

This leads to a disciplined viewpoint:

- "Evaluation at infinity" is a local problem in the infinity chart.
- Under w = 1/n, it becomes "evaluation near a puncture at w = 0."
- In either chart, a directional limit requires a direction or path, because the function's asymptotics vary across sectors.

Thus the coordinate change w = 1/n is not merely a convenience. It is a conceptual proof that the gamma-data is not an afterthought. The direction is intrinsic to the phenomenon. Infinity is a single point, but the analytic neighborhood of that point has a rich angular structure, and our notation n_infty paired with gamma_theta is precisely a way of recording which angular sector we inhabit when we claim that an exponential "vanishes at infinity."

**7. A Minimal Calculus of Directional Infinity**

Once we allow directional evaluation at n_infty, the next question is how far we can push it without accidentally smuggling in contradictions. The goal here is modest: we do not want to build a new number system or a full algebra of infinity. We want a minimal calculus that tells us when f(n_infty; gamma) exists, how to compute it for common classes of functions, and what operations are safe. The guiding principle is simple: n_infty is not a number; f(n_infty; gamma) is a limit. Limits can be manipulated under familiar hypotheses, but the hypotheses must be stated. When those hypotheses are ignored, "infinity arithmetic" appears to work until it suddenly yields 1 = 0. This section collects the basic rules in one place.

**7.1 Existence criteria for f(n_infty; gamma) for common function classes**

Recall the definition: for a curve gamma(t) in C with |gamma(t)| -> infinity as t -> +infty, define

f(n_infty; gamma) = lim(t -> +infty) f(gamma(t))

when the limit exists.

So existence is always a question about an ordinary limit along a path. There is no universal guarantee. Different function classes behave differently, but a useful starting heuristic is:

- If f is "asymptotically constant" along gamma, the limit exists and equals that constant.
- If f has growth, blow-up, or oscillation along gamma, the limit may fail to exist or may be infinite.
- If f is meromorphic (rational functions are the main example), infinity behaves like another point where poles and zeros can occur, and the limit along any path is often computable.
- If f involves exponentials, existence is typically governed by the sign of the real part of the exponent along gamma.

In practice, the simplest existence check is to parameterize the chosen gamma and analyze f(gamma(t)) as t -> +infty, exactly as we did for exp(i pi theta gamma_theta(t)).

Two levels of “existence” are worth distinguishing.

Directional existence: the limit exists along the specific gamma.
Global existence: the limit exists and is the same along all paths to infinity, so f extends continuously to n_infty on the Riemann sphere.

Our framework is primarily directional. Global existence is stronger and rarer; it holds for functions like 1/n, but fails for many exponentials and oscillatory expressions.

**7.2 Rational functions: poles, finite limits, and divergence at n_infty**
Rational functions provide the cleanest testing ground because their behavior at infinity is completely controlled by degrees.

Let r(n) = p(n)/q(n), where p and q are polynomials with no common factor and q not identically zero. Then as |n| -> infinity, r(n) behaves like the ratio of leading terms. That immediately yields:

- If deg p < deg q, then r(n) -> 0 as n -> n_infty, and the limit is direction-independent. In our notation, r(n_infty; gamma) = 0 for every gamma.
- If deg p = deg q, then r(n) -> (leading coefficient of p) / (leading coefficient of q), again direction-independent. So r has a finite “value at infinity” in the global sense.
- If deg p > deg q, then |r(n)| -> infinity as n -> n_infty along any path that avoids zeros of q, and there is no finite limit. In meromorphic language, r has a pole at infinity of order deg p - deg q.

This is the first place where “infinity” behaves exactly like an ordinary point in function theory. Saying “r has a pole at infinity” is not metaphor. It is a precise statement about r as a meromorphic function on CP1.

Directional subtleties for rationals are relatively mild. You can still have path-dependent behavior if you insist on approaching infinity along curves that repeatedly pass near poles or zeros in a contrived way, but for typical rays and for degree arguments, the asymptotics are stable and direction-independent. This makes rational functions a useful baseline: they show that n_infty is not inherently problematic; the difficulties arise when functions have essential direction-sensitive growth or oscillation near infinity.

**7.3 Exponentials and exponential polynomials: dominance by real parts**
Exponentials introduce the main new feature: sectorial dominance governed by real parts.

Start with the simplest case: f(n) = exp(a n), with a a fixed complex constant. Along a ray n = r u with r -> +infty and |u| = 1,

|exp(a n)| = exp( Re(a r u) ) = exp( r Re(a u) ).

So:

- If Re(a u) < 0, then exp(a n) -> 0 along that ray.
- If Re(a u) > 0, then exp(a n) blows up along that ray.
- If Re(a u) = 0, then |exp(a n)| = 1 along that ray and the behavior is oscillatory; a limit typically does not exist unless the phase stabilizes in some special way.

This is exactly the mechanism behind our construction with a = i pi theta and u chosen so that a u is a negative real number.

Now consider exponential polynomials, finite sums of exponentials with polynomial coefficients:

f(n) = sum_{k=1}^m P_k(n) exp(a_k n),

where each P_k is a polynomial and each a_k is a complex constant. Along a fixed ray n = r u, the magnitudes behave like

|P_k(r u) exp(a_k r u)| approx (polynomial growth in r) times exp(r Re(a_k u)).

As r -> +infty, the exponential factors dominate the polynomial ones. Therefore, along that ray, the term(s) with the largest value of Re(a_k u) dominate the asymptotics.

- If the maximum Re(a_k u) is negative and achieved uniquely, then f(n) -> 0 along that ray, because every term decays exponentially and polynomial factors cannot prevent decay.
- If the maximum is positive, then f grows without bound in magnitude along that ray unless there is an exact cancellation between equally dominant terms, which is nongeneric and unstable under perturbation.
- If the maximum is zero, then the dominant terms are oscillatory in magnitude, and f typically does not converge to a limit; it may remain bounded, oscillate, or fail to settle.

This “dominance by real parts” is the seed of the sector picture used throughout asymptotics: different rays u pick out different dominant exponentials by changing the ordering of Re(a_k u). Boundary rays where two exponents tie in real part are precisely where dominance switches, and where asymptotic expansions often require special handling.

For our single exponential exp(i pi theta n), there is only one exponent, so the story is simpler: existence of a limit is governed entirely by whether Re(i pi theta u) is negative, positive, or zero along the chosen direction u.

**7.4 Warnings: why naive algebra with n_infty produces contradictions**

The notation n_infty is geometrically meaningful, but it is not a field element. The boundary functional f(n_infty; gamma) is a limit, and limits do not automatically respect all algebraic manipulations unless the relevant limits exist and the operations are continuous in the required sense.

The quickest way to see the danger is to notice how easy it is to create undefined forms. For example, if you try to treat 1/n_infty as a number, you will be tempted to write 1/n_infty = 0. That is often a correct limit statement in the sense that (1/n)(n_infty; gamma) = 0 for many gamma. But if you then combine it with n_infty * (1/n_infty) = 1, you are mixing limit-based interpretation with algebraic interpretation. The expression n_infty * 0 is not defined in a way that preserves the field axioms, and if you pretend it is, you will quickly obtain contradictions.

The safe rule is:

- Replace every expression involving n_infty with the corresponding limit along a stated gamma, and only then perform algebraic manipulations that are justified for limits.

For instance, it is safe to say:
If lim f(gamma(t)) = A and lim g(gamma(t)) = B and both limits are finite, then lim (f+g)(gamma(t)) = A+B and lim (f g)(gamma(t)) = A B. This is ordinary limit algebra.

But it is not safe to say:
If f(n_infty; gamma) = A, then f(n_infty) = A as an unconditional value. That forgets gamma, and it may be false.

It is also not safe to manipulate expressions that produce indeterminate forms:
infinity minus infinity, 0 times infinity, or exp(infinity) without specifying whether the real part tends to plus or minus infinity.

In particular, returning to the original motivating temptation, it is never valid to reason:
exp(i pi theta n_infty) = 0 therefore exp(-i pi theta n_infty) is its inverse. The inverse law exp(z) exp(-z) = 1 holds for finite z. At the boundary, exp(i pi theta n_infty; gamma_theta) = 0 is a limit statement, and multiplying by another limit that diverges can yield an indeterminate form rather than 1. The formal identity does not transport to the boundary without extra hypotheses.

So the minimal calculus is intentionally conservative:

- n_infty is an ideal point.
- gamma records approach.
- f(n_infty; gamma) is a limit when it exists.

- Algebra is done at the level of limits, not by treating infinity as a number.

Within those constraints, the framework remains rigorous, expressive, and free of contradiction, while still allowing the concise statement that exp(i pi theta n) “vanishes at directional infinity” along gamma_theta._

**8. Applications as Boundary Conditions and Selection Rules**

Directional infinity is not merely a linguistic repair for a paradoxical-looking equation. Its real value is that it upgrades a common informal move into a precise, reusable selection principle: impose a boundary condition “at infinity,” and you can pick one solution among many. This is everywhere in analysis. The difficulty is that in complex settings “at infinity” is ambiguous unless a direction is specified. Once we name n_infty and attach an approach ray gamma, the boundary condition becomes an ordinary limit constraint. In this section we show how the condition f(n) -> 0 as n -> n_infty along gamma_theta functions as a clean selection rule; how it fits naturally into ordinary differential equations and discrete recurrences; how it provides a geometric interpretation of damping and regularization; and how it parallels the logic behind steepest descent, where contour choice is exactly the choice of decay direction.

**8.1 Using f(n) -> 0 as n -> n_infty along gamma_theta to pick solutions**

Many problems do not have a unique solution until one adds an asymptotic requirement. In real-variable settings this is familiar: two linearly independent solutions exist to a second-order ODE, and one chooses the “decaying” one by imposing a boundary condition at infinity. In complex-variable settings the same logic holds, but the phrase “decaying at infinity” must be interpreted directionally.

The directional boundary condition takes the form:

Choose the solution f such that lim(t -> +infty) f(gamma_theta(t)) = 0.

This is a single scalar condition on the behavior of f along a specified ray. Yet it often has strong consequences. For linear problems, it typically eliminates an entire subspace of solutions, leaving a one-dimensional space or a unique solution after normalization.

A simple archetype is any equation whose solutions include terms of the form exp(a n). If two candidate solutions differ by a term that grows along gamma_theta, then the decay condition rules that candidate out. In this way, gamma_theta becomes a “filter” that selects the solution compatible with the decay half-plane determined by the exponent coefficient.

In our setting, gamma_theta is engineered to make exp(i pi theta n) decay. So a boundary condition “vanish along gamma_theta” is an instruction that the solution must live in the

analytic sector where i pi theta n has large negative real part. This is exactly the kind of instruction that turns an underdetermined problem into a determined one.

The conceptual point is that the selection rule is not mystical. It is simply the imposition of a directional limit at a boundary point of the domain. But because complex infinity is a single point approached by many rays, the direction is what makes the selection rule meaningful rather than contradictory or vacuous.

**8.2 A template for ODE and recurrence problems with "decay at infinity" conditions**

The same selection idea appears in two broad families of problems: continuous differential equations and discrete recurrences. The directional infinity framework gives a standard template for stating and using the boundary condition.

Template for ODE problems.
Consider a linear ODE in a complex variable n (or z), for example

Lf = g(n),

where L is a linear differential operator with analytic coefficients on a region of C. The solution space of the homogeneous equation L[f] = 0 typically has dimension equal to the order of L, and one needs extra conditions to isolate a particular solution. The directional infinity condition is:

Impose f(n) -> 0 as n -> n_infty along gamma, for a chosen gamma.

Operationally, one proceeds as follows:

- Identify the asymptotic modes of solutions (often exponential-like).
- Determine which modes decay along the chosen ray gamma by examining real parts of exponents.
- Set the coefficients of the growing modes to zero.
- Solve for the remaining coefficients using initial conditions or normalization constraints.

This is precisely how one selects Airy-type or WKB-type solutions in classical asymptotics: one specifies a sector (a direction at infinity) in which the solution should decay, which picks out a canonical branch.

Template for recurrence problems.
Now consider a linear recurrence in an index n, extended to complex n when appropriate, or simply analyzed asymptotically as n grows large:

A(n) f(n+1) + B(n) f(n) + C(n) f(n-1) = h(n),

with coefficients A, B, C that may have asymptotic expansions. Such recurrences often have multiple asymptotic behaviors: one solution may grow, another may decay, another may oscillate. In discrete problems the phrase “as n -> infinity” usually means n along the positive integers, but the same conceptual ambiguity can appear if one analytically continues n or studies generating functions where complex directions matter.

Even without full continuation, the spirit of directional selection is still useful: decide which asymptotic mode is admissible (decay vs growth), and discard the others. In the discrete setting, the analog of “choose gamma” is “choose the asymptotic regime” (for example, the mode that decays for large positive integer n). When a complexified viewpoint is appropriate, gamma becomes literal again: one studies f(gamma(t)) for a curve gamma in C and imposes decay along that curve.

So the framework offers a uniform language: whether continuous or discrete, a “decay at infinity” condition is a selection rule on asymptotic modes, and in complex settings it must always be tied to a sector or ray.

**8.3 Regularization intuition: damping as a geometric choice of direction**

A common practical move in analysis is to “insert damping” to force convergence. One might multiply an oscillatory integrand by exp(-epsilon t) and later take epsilon -> 0, or one might tilt a contour slightly into the complex plane so that exp(i t) becomes exp(i t - epsilon t). Directional infinity provides a geometric interpretation of this move: damping is often nothing more than choosing a direction in which the real part of an exponent becomes negative.

In our construction, exp(i pi theta n) appears purely oscillatory when theta and n are real. The moment we allow n to acquire an imaginary component in the correct sign, the exponent gains a negative real part and the oscillation becomes exponentially damped. The choice of gamma_theta is exactly the “damped direction” written geometrically. It is the cleanest example of a general phenomenon:

- Oscillatory terms become decaying terms by moving into a complex direction where the exponent’s real part is negative and large in magnitude.

From this viewpoint, regularization is not always an external modification of a function. It can be an internal geometric choice of how we probe the function near infinity. Instead of adding epsilon by hand, we select a path gamma that naturally supplies the negative real part that epsilon would have introduced.

This is especially clarifying when discussing boundary values, Fourier-type oscillations, and analytic continuation. Many apparent divergences are not absolute; they are artifacts of staying on a boundary ray where the real part vanishes. A small tilt into a decay sector is equivalent to

choosing a gamma that crosses into Re(exponent) < 0 territory. The framework packages that tilt as a directional boundary condition, not as an ad hoc damping term.

**8.4 Relation to steepest descent contours and "best" decay directions**

The deepest conceptual neighbor of our construction is steepest descent. In the method of steepest descent, one evaluates integrals of the form

Integral exp(lambda Phi(z)) A(z) dz

for large lambda by deforming the contour into the complex plane so that Re(lambda Phi(z)) decreases as rapidly as possible away from critical points. The method is fundamentally about choosing paths along which exponentials decay. The contour is chosen so that oscillation becomes decay, allowing the integral to be dominated by neighborhoods where the phase is stationary.

Our ray gamma_theta is the simplest steepest descent contour imaginable: Phi(n) = i pi theta n is linear, so there are no critical points to locate; the entire problem is to choose a direction in which Re(Phi(n)) goes to minus infinity. The "best" decay direction is the one that makes Phi(n) purely real and negative, because then Re(Phi(n)) is as negative as possible for a given |n|. That is precisely what gamma_theta does by construction: it makes i pi theta gamma_theta(t) equal to -t.

So gamma_theta can be read as a steepest descent path for a linear phase. The sector picture from earlier sections is the same geometry that underlies steepest descent: decay regions are sectors where Re(Phi) is negative, boundary rays are where Re(Phi) is zero, and growth regions are where Re(Phi) is positive.

This connection matters because it explains why the framework is not a one-off invention. The idea "choose a path so the exponential decays" is a central organizing principle of complex asymptotics. Directional infinity is a compact way to state the same principle as a boundary condition at n_infty: among all possible approaches to infinity, select those consistent with decay, and treat that selection as part of the problem data.

In that sense, the expression exp(i pi theta n_infty; gamma_theta) = 0 is not an exotic claim. It is the boundary-value shorthand for "we are in the steepest descent sector." It formalizes the intuitive practice of analysts and asymptoticians: choose the direction that makes the exponential vanish, and then build the rest of the argument inside that decay sector.

**9. Extensions and Variants**

The construction so far is intentionally minimal: a single ideal point n_infty, a single preferred ray gamma_theta, and a single boundary functional f(n_infty; gamma). But the underlying ideas

are more general. Infinity in complex analysis is simultaneously simple (a single added point in CP1) and rich (many distinct approach directions with distinct asymptotics). Once we accept that direction is part of the data, there are several natural directions to extend the framework: parameterize whole families of approach rays, encode direction more intrinsically by refining the boundary geometry, generalize from linear exponents to broader exponential classes, and lift the entire picture from one complex variable to several, where “direction” becomes a cone rather than a single angle.

**9.1 Families of rays gamma(theta, phi) and directional parameterization**

For a fixed theta != 0, we selected one canonical ray gamma_theta(t) that forces i pi theta gamma_theta(t) = -t. That ray is convenient because the proof becomes a one-line reduction, and because it is a “maximal decay” direction among straight rays. But there is not only one decay ray. There is an entire half-plane of directions along which the exponential decays, and within that half-plane one may want to choose different rays for different analytic tasks.

A natural way to describe a family is to parameterize rays by an angle phi:

gamma(phi; t) = t exp(i phi), t > 0.

Then the decay condition for exp(i pi theta n) along gamma(phi; t) becomes a condition on phi:

|exp(i pi theta t exp(i phi))| = exp( t Re(i pi theta exp(i phi)) ).

Decay occurs when Re(i pi theta exp(i phi)) < 0.

For a fixed theta, this inequality defines an open interval of angles phi, a “decay arc” on the unit circle. The endpoints of that interval are boundary angles where Re(i pi theta exp(i phi)) = 0, which correspond to the boundary rays separating decay from growth. In this parameterization, gamma_theta is just one selected phi inside the decay interval, chosen so that i pi theta exp(i phi) is a negative real number rather than merely having negative real part.

If one wants an explicit two-parameter family that includes both the theta-dependence and the freedom within the decay half-plane, one can define gamma(theta, s) by requiring

i pi theta gamma(theta, s; t) = -(1 + s i) t,

where s is a real parameter. This yields

gamma(theta, s; t) = - (1 + s i) t / (i pi theta).

For any fixed s, the real part of the exponent is -t, so the magnitude decays like exp(-t) while the imaginary part produces additional oscillation. This family shows that there are infinitely many “equally decaying” rays (same exponential rate) differing only in the oscillatory component. The original gamma_theta corresponds to s = 0, the purely real exponent case.

This kind of parameterization is practical: it separates the decay rate (set by the real part) from oscillation (set by the imaginary part), and it makes the role of “directional choice” explicit as a tunable parameter rather than a single fixed path.

**9.2 Encoding direction intrinsically: blow-up of infinity and a boundary circle of directions**
The Riemann sphere adds a single point at infinity, which is topologically and algebraically elegant, but it collapses all approach directions into one location. When direction is essential, it is natural to refine the compactification so that different approach directions become distinct boundary points.

One standard geometric refinement is to “blow up” the point at infinity. Informally, this replaces the single point n_infty with a circle (or projective line) of directions. The circle parametrizes approach angles: each point on the circle corresponds to a direction in which one can escape to infinity. Under this refinement, the pair (n_infty, gamma) becomes a single boundary point: infinity together with its approach direction.

In one complex variable, the space of directions to infinity can be identified with the unit circle modulo antipodal identification, depending on whether one considers oriented rays or unoriented lines. If we care about rays (oriented), the direction space is essentially the unit circle. If we care only about lines through the origin, it is the circle with antipodal points identified.

Why is this useful? Because it turns “directional evaluation at infinity” into ordinary evaluation on a larger boundary. Instead of writing f(n_infty; gamma), one can speak of f at a boundary point labeled by an angle. This makes sector analysis feel intrinsic: decay sectors become arcs of the boundary circle; boundary rays become boundary points separating those arcs.

Even if one does not implement the full blow-up geometry formally, this perspective is conceptually valuable. It makes clear that direction is not an auxiliary annotation; it is missing boundary structure that can be restored by refining the compactification.

**9.3 Beyond exp(i pi theta n): general exp(a n) and exp(p(n)) cases**
The example exp(i pi theta n) is a prototype for exp(a n) with a a fixed complex constant. Everything we did generalizes immediately by replacing a = i pi theta:

- Choose a ray gamma_a(t) such that a gamma_a(t) = -t.
- Then exp(a gamma_a(t)) = exp(-t) -> 0.

Explicitly, for any a != 0 one can define

gamma_a(t) = -t / a,

and obtain the same one-line decay proof. So the entire framework is really about exponentials with linear phase. The special form i pi theta is not essential; it simply emphasizes the oscillatory appearance when theta is real.

A more interesting generalization is exp(p(n)) where p is a polynomial (or more broadly, an analytic function) with leading term controlling behavior at infinity. Here the geometry becomes richer because the sign of Re(p(n)) can vary by direction in a more intricate way. For example, if p(n) = n^2, then

Re(n^2) = r^2 cos(2 phi)

along n = r exp(i phi). This produces four alternating sectors of decay and growth for exp(-n^2) or exp(n^2), separated by boundary rays where cos(2 phi) = 0. This is the simplest example of the multi-sector phenomenon that underlies Stokes lines and steepest descent in non-linear phases.

In general, for p(n) with leading term c n^m, the dominant behavior for large |n| along a ray depends on Re(c exp(i m phi)) times r^m. This yields m-fold sectoring: the plane divides into 2m alternating sectors where Re(c n^m) is positive or negative, and those sectors control decay or growth of exp(p(n)).

So the “direction matters” principle scales naturally:

- For linear phases, there are two main half-planes (decay and growth) separated by two boundary rays.
- For higher-degree phases, there are multiple alternating decay and growth sectors, and the choice of ray becomes even more consequential.

In such cases, a natural analogue of gamma_theta is not necessarily a straight ray, but a steepest descent curve defined by keeping Im(p(n)) constant while forcing Re(p(n)) to decrease most rapidly. Directional infinity still makes sense, but the preferred paths may be curved rather than straight, and the boundary data becomes “approach within a chosen sector” rather than “approach along this single ray.”

**9.4 Multivariable generalization: infinity in C^k and approach cones**

When n is replaced by a vector z in C^k, the geometry of infinity becomes higher-dimensional. There are multiple inequivalent ways to compactify C^k, and the space of “directions to infinity” is no longer a circle but a higher-dimensional sphere-like object. The basic phenomenon, however, remains: different directions to infinity can produce different asymptotics, and for exponentials the sign of a real part controls decay.

For linear exponentials in several variables, consider

f(z) = exp( <a, z> ),

where a and z are in C^k and <a, z> is a complex bilinear pairing. If we take z = r u with r -> +infty and u a direction vector, then

|f(r u)| = exp( r Re(<a, u>) ).

Decay occurs in directions u for which Re(<a, u>) < 0. This defines a half-space in the space of directions. So in C^k, the set of decay directions is a region on the unit sphere of directions; growth directions form the complementary region; and the boundary where Re(<a, u>) = 0 is a codimension-one “great hypersurface” separating them.

In this setting, it is more natural to speak of approach cones rather than rays. A cone is a set of directions rather than a single direction: for example, all u in a neighborhood on the direction sphere such that Re(<a, u>) stays negative. If a path to infinity eventually stays inside such a cone, then f(z) decays along that path.

For polynomial phases p(z) in several variables, the situation becomes richer again: the sign of Re(p(r u)) is controlled by the leading homogeneous part of p, and the direction sphere is partitioned into multiple regions where Re(p) tends to plus or minus infinity at different rates. This leads to a higher-dimensional analogue of sectoring and Stokes phenomena.

So the key upgrades in C^k are:

- Direction data is no longer one angle; it is a point (or region) on a higher-dimensional direction sphere.
- The natural “directional infinity” object is a cone (or a collection of cones) rather than a single ray.
- The boundary functional becomes f(infty; cone) meaning a limit along any path escaping to infinity within a specified cone, when such a limit is stable across that cone.

These multivariable generalizations show that the framework is not tied to one-dimensional quirks. It is an instance of a general pattern: compactify the domain, treat infinity as boundary, and recognize that asymptotics live on the boundary together with direction. The one-variable symbol n_infty and the ray gamma_theta are simply the simplest nontrivial case where this pattern resolves an apparent paradox cleanly.

## 10. Conclusion

The thread that motivated this paper began with a seemingly impossible request: choose an index n so that exp(i pi theta n) equals zero. In ordinary complex analysis that request has no solution, and the impossibility is not technical but structural: the exponential never vanishes. The work of the paper has been to preserve that structural fact while still capturing the intuition hiding behind the request, namely the desire to talk about “vanishing behavior” in a clean, symbolic way. The resolution is to move from value language to boundary language. Instead of inventing a forbidden n, we name the ideal point at infinity n_infty on the Riemann sphere, and we attach an approach ray gamma_theta along which the exponent becomes real and tends to minus infinity. Along that ray, the exponential decays to zero in the only sense compatible with exp’s nonvanishing: as a directional limit. What looks like a paradox is revealed to be a mismatch of grammars: pointwise equality versus asymptotic boundary evaluation.

### 10.1 The resolution: no contradiction with exp(z) != 0

The nonvanishing theorem remains untouched throughout. For every finite complex z, exp(z) is nonzero and has an inverse exp(-z). The expression exp(i pi theta n) therefore cannot equal zero for any finite n. The paper’s central move is that it never tries to force such an equality.

Instead, it introduces a boundary functional: for a path gamma(t) with |gamma(t)| -> infinity, the statement

f(n_infty; gamma) = L

means lim(t -> +infty) f(gamma(t)) = L. When f(n) = exp(i pi theta n) and gamma = gamma_theta(t) = -t/(i pi theta), we obtain

exp(i pi theta n_infty; gamma_theta) = 0

because the exponent equals -t along that ray. This is a limit statement, not a value statement. There is no z with exp(z) = 0; there is only a path to infinity along which exp(z) tends to 0. The distinction is not philosophical hair-splitting; it is exactly what prevents contradictions like 1 = 0. The exponential’s algebraic identities live on the finite domain; the boundary evaluation lives in a topology of limits, where indeterminate forms can arise unless handled with care.

### 10.2 Infinity as geometry plus approach data, not a magical value

Infinity becomes mathematically usable when treated as geometry, not arithmetic. On the Riemann sphere (or CP1), infinity is a single added point that compactifies the complex plane. This makes statements “at infinity” analogous to statements at ordinary points: we can discuss limits, continuity, and extension of functions to the boundary.

But complex analysis adds a second layer: the point at infinity does not remember the direction of approach. Many functions, especially exponentials and compositions involving poles, have

different limiting behavior along different rays that all converge to the same n_infty. That is why the approach path gamma is not an afterthought. It is part of the meaning of an asymptotic claim.

The pair (n_infty, gamma) is therefore the correct conceptual object: infinity provides the boundary address, and gamma provides the analytic content. The coordinate change w = 1/n reinforces this: “approach infinity” becomes “approach the puncture w = 0,” and the same directional dependence reappears as the direction of approach to the puncture. Direction is not a quirk of infinity. It is an intrinsic feature of local complex behavior near boundary or singular points.

**10.3 A compact language for vanishing, growth, and oscillation at the boundary**

Once direction is treated as data, the asymptotic trichotomy for exponentials becomes clean and computable. For exp(a n), behavior along a ray is controlled by the sign of Re(a u), where u is the ray direction:

- Negative real part yields decay to 0.
- Positive real part yields growth to infinity in magnitude.
- Zero real part yields unit-magnitude oscillation and typically no limit.

This can be expressed either as a sector decomposition in the n-plane or as a directional boundary evaluation at n_infty. The notation f(n_infty; gamma) packages the limit into a single symbol without smuggling in false equalities. It also makes it easy to state selection rules: choose the solution that vanishes along a specified ray, or restrict attention to paths lying in a decay half-plane or decay sector.

In that way, the framework provides a compact language that mirrors what analysts actually do: they choose directions (or contours) to force decay, they track boundary rays where behavior changes, and they interpret “vanishing at infinity” as a controlled limit rather than as a literal value.

**10.4 What the framework buys: rigor, reuse, and clean asymptotic reasoning**

The payoff is practical.

Rigor: The framework replaces an impossible equation with a precise, well-posed limit statement. It preserves the nonvanishing of exp while still allowing one to speak meaningfully about “exponential vanishing” in the only consistent sense: decay along an approach to the boundary.

Reuse: The same structure applies immediately to exp(a n) for any a != 0, and extends naturally to exp(p(n)) where p is polynomial, producing the familiar sectoring patterns of higher-degree

phases. It also lifts to several complex variables, where rays generalize to cones of approach directions.

Clean reasoning: By forcing the approach path to be named, the framework prevents a common source of confusion: claiming a limit “at infinity” without specifying whether it is unconditional or directional. It clarifies why some limits are direction-independent (as with 1/n) while others are inherently sectorial (as with exponentials). It aligns naturally with standard methods such as steepest descent, where contour choice is the analytic mechanism that turns oscillation into decay.

In short, what began as a request for a special n becomes a small but powerful reframing: do not ask for a value that violates the function’s algebra; ask for a boundary direction that reveals the function’s asymptotics. Infinity is not a number you solve for. It is a point you approach, and the approach carries the meaning.

## 11. References


[1] Lars V. Ahlfors. Complex Analysis (Ahlfors): An Introduction to the Theory of Analytic Functions of One Complex Variable. 3rd ed. McGraw-Hill, 1979. ISBN 978-0-07-000657-7.

[2] John B. Conway. Functions of One Complex Variable I. 2nd ed. Graduate Texts in Mathematics 11. Springer-Verlag, 1978. ISBN 978-0-387-90328-6.

[3] Elias M. Stein and Rami Shakarchi. Complex Analysis (Stein–Shakarchi). Princeton Lectures in Analysis, Vol. II. Princeton University Press, 2003. ISBN 978-0-691-11385-2.

[4] Tristan Needham. Visual Complex Analysis. Oxford University Press, 1997. ISBN 0-19-853446-9.

[5] Theodore W. Gamelin. Complex Analysis (Gamelin). Undergraduate Texts in Mathematics. Springer, 2001. ISBN 0-387-95093-1.

[6] E. C. Titchmarsh. The Theory of Functions. 2nd ed. Oxford University Press, 1939.

[7] Otto Forster. Lectures on Riemann Surfaces. Graduate Texts in Mathematics 81. Springer, 1981. ISBN 978-0-387-90617-1.

[8] Hershel M. Farkas and Irwin Kra. Riemann Surfaces (Farkas–Kra). 2nd ed. Graduate Texts in Mathematics 71. Springer, 1992. ISBN 978-0-387-97703-4.

[9] Rick Miranda. Algebraic Curves and Riemann Surfaces. Graduate Studies in Mathematics 5. American Mathematical Society, 1995. ISBN 978-0-8218-0268-7.

[10] Robin Hartshorne. Algebraic Geometry. Graduate Texts in Mathematics 52. Springer, 1977. ISBN 978-0-387-90244-9.

[11] Phillip Griffiths and Joseph Harris. Principles of Algebraic Geometry. John Wiley & Sons, 1978. ISBN 978-0-471-05059-9.

[12] John Milnor. Dynamics in One Complex Variable. 3rd ed. Princeton University Press, 2006.

[13] Frank W. J. Olver. Asymptotics and Special Functions. Academic Press, 1974. ISBN 978-0-12-525850-0.

[14] Roderick Wong. Asymptotic Approximations of Integrals. SIAM, 2001. ISBN 978-0-89871-497-5.

[15] Norman Bleistein and Richard A. Handelsman. Asymptotic Expansions of Integrals. Dover Publications, 1986. ISBN 978-0-486-65082-1.

[16] R. B. Dingle. Asymptotic Expansions: Their Derivation and Interpretation. Academic Press, 1973. ISBN 978-0-12-216550-4.

[17] R. B. Paris and David Kaminski. Asymptotics and Mellin-Barnes Integrals. Cambridge University Press, 2001. ISBN 978-0-521-79463-3.

[18] M. V. Berry. “Uniform asymptotic smoothing of Stokes’s discontinuities.” Proceedings of the Royal Society of London. Series A, 422 (1989): 7–21.

[19] N. G. de Bruijn. Asymptotic Methods in Analysis. Dover, 1981 (originally published 1958).

[20] G. H. Hardy. Divergent Series. Oxford University Press, 1949.

[21] Ralph P. Boas Jr.. Entire Functions. Academic Press, 1954.

[22] B. Ya. Levin. Distribution of Zeros of Entire Functions. American Mathematical Society (English translation), 1980; original Russian edition 1964.

[23] Carl M. Bender and Steven A. Orszag. Advanced Mathematical Methods for Scientists and Engineers: Asymptotic Methods and Perturbation Theory. Springer, 1999. ISBN 978-0-387-98931-2.

The author has published ~ 5,000 AI-assisted papers at: https://www.researchgate.net/profile/Douglas-Youvan/research . Google Scholar has indexed these preprints, and if you want a particular reference, the AI mode of a Google search (Gemini) has efficiently catalogued these preprints.


∞ (n∞)
Decaying
exp(iπ θ1 n)
?θ1
Growing
exp(iπ θ2 n)
w = r/in
w → 0
Exponential
Decay → 0
n → ∞
w → 0
Oscillation Re(iπθn)